

Primary Author: Hemanth Venkatesh, MD

Editor: Michael L. Cheatham MD, Chadwick P. Smith, MD

Approved: 04/01/2004

Revised 10/27/2009, 10/30/2024

## SUMMARY

Pulmonary embolism (PE) remains a significant cause of morbidity and mortality in the critically ill surgical or trauma patient. PE may occur even in the presence of appropriate deep venous thrombosis (DVT) prophylaxis. Patients at high risk for PE may benefit from placement of an inferior vena cava (IVC) filter if they cannot be anticoagulated. While these devices have been shown to be effective in the prevention of PE, they are associated with an increased risk of DVT and have not been proven to reduce mortality.

## RECOMMENDATIONS

### Level 1

- **Routine prophylactic IVC filter insertion in trauma patients with ISS < 15 should not be performed.**
- **Routine IVC filter placement is not indicated in patients with DVT who can be anticoagulated.**

### Level 2

- **IVC filter insertion is indicated in patients with proximal DVT who cannot be anticoagulated. Such patients should be anticoagulated when their bleeding risk resolves.**
- **Temporary IVC filters may be considered when the risk of PE or contraindications to anticoagulation is anticipated to be less than two (2) weeks and the risk of PE is high.**
- **IVC filters may be safely placed at the patient's bedside under either fluoroscopic or ultrasound guidance.**

### Level 3

- **None**

## INTRODUCTION

Deep venous thrombosis (DVT) and pulmonary embolism (PE) remain common, challenging, and often-devastating complications in the surgical or trauma patient. The average incidence of DVT in the general trauma population is 42% (range 18-90%) and the reported incidence of PE in patients with spinal cord injury (SCI) is 10% (range 4%-22%). Up to 4% of injury-related deaths in the U.S. are caused by PE-related "sudden death", frequently in patients that would otherwise have recovered from their injuries. A patient's risk increases within the first several hours after injury with DVT and/or PE frequently being noted within the first 72 hours. Reports exist of PE in the first 24-48 hours post-injury.

PE following development of DVT is one of the most preventable causes of death in hospitalized patients. DVT prophylaxis using either unfractionated / fractionated heparin or intermittent pneumatic compression devices represents the first-line of therapy but is neither 100% protective against DVT formation nor subsequent PE. This is especially true in the critically ill, high-risk patient who may have barriers or contraindications to the use of such methods of prophylaxis such as complex wounds, CNS (brain and spinal cord) or ocular injuries, external fixators, or traction devices.

IVC filters have been proven to decrease the risk of PE in various patient populations including the critically ill and traumatically injured. Reported complication rates range from 0-35% with patency rates in excess of 90%. Concerns

## LEVEL OF RECOMMENDATION DEFINITIONS

- **Level 1:** Supported by multiple, prospective randomized clinical trials or strong prospective, non-randomized evidence if randomized testing is inappropriate.
- **Level 2:** Supported by prospective data or a preponderance of strong retrospective evidence.
- **Level 3:** Supported by retrospective data or expert opinion.

DISCLAIMER: These guidelines were prepared by the Department of Surgical Education, Orlando Regional Medical Center. They are intended to serve as a general statement regarding appropriate patient care practices based on the medical literature and clinical expertise at the time of development. They should not be considered to be accepted protocol or policy, nor are intended to replace clinical judgment or dictate care of individual patients.

include the safety and long-term effects of these devices, especially in younger patients, for whom the risk of thromboembolism may be time-limited. The recent availability of removable devices may solve some of these problems, offering protection against PE during the early, highest-risk period, while avoiding the potential long-term complications of a permanent filter. To-date, however, few studies have shown that these filters are truly “temporary” with many such devices being left in place permanently.

The Eastern Association for the Surgery of Trauma (EAST) has published extensive evidence-based medicine guidelines on the management of DVT in the trauma patient (1). These guidelines, which have not been updated since 2002, recommend IVC filter placement in patients with the following findings:

- Recurrent PE despite full anticoagulation (Level I)
- Proximal DVT and contraindications to full anticoagulation (Level I)
- Proximal DVT and major bleeding while on full anticoagulation (Level I)
- Progression of iliofemoral clot despite anticoagulation (rare) (Level I)
- Large free-floating thrombus in the iliac vein or IVC (Level II)
- Following massive PE in which recurrent emboli may prove fatal (Level II)
- During/after surgical embolectomy (Level II)
- “Prophylactic” vena cava filter insertion in very high-risk trauma patients who: (Level III)
  1. Cannot receive anticoagulation because of increased bleeding risk, and
  2. Have one or more of the following injury patterns:
    - Severe closed head injury (GCS < 8)
    - Incomplete spinal cord injury with para or quadriplegia
    - Complex pelvic fractures with associated long-bone fractures
    - Multiple long-bone fractures

The American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (9th Edition) recommends the following regarding IVC filter placement (2):

- In patients with acute DVT of the leg, we recommend against the use of IVC filter in addition to anticoagulants (strong recommendation, moderate-certainty evidence).
- In patients with acute proximal DVT of the leg and a contraindication to anticoagulation, we recommend the use of an IVC filter (strong recommendation, moderate-certainty evidence).

The American College of Surgery (ACS) recently published a ‘Best Practice Guidelines’ on ‘The Management of Traumatic Brain Injury’. In this, they discuss the indication for IVC filters being the presence of a known VTE in a patient with a contraindication for therapeutic anticoagulation. They explicitly state “use of prophylactic inferior vena cava infilters is no longer recommended, regardless of the timing of pharmacologic VTE prophylaxis in TBI patients.” (3).

Despite the above practice guideline from ACS, others have demonstrated a mortality benefit to IVC filters in the severely injured trauma patients with a neurotrauma component. Elkbuli et. Al. performed a retrospective review of trauma patients > 16 years of age with ISS > 15. 413 patients received prophylactic IVC filters. Patients with IVC filters prophylactically placed had higher ISS, lower GCS, and higher prevalence of TBI. Patients in the IVC filter group had higher rates of DVT and nonfatal PE along with a prolonged ICU length of stay. Of note, among patients with a head injury score > 3, prophylactic IVC filter placement was associated with lower in hospital mortality compared to VTE prophylaxis (4). All in all, this paper highlights the importance of further research into evaluating the mortality benefit of neurotrauma patients.

**LITERATURE REVIEW**

**Indications for IVC Filter Insertion**

Author	Year	Evidence	Findings
Leach (5)	1985	Class II	Review of 201 trauma patients. Negligible morbidity and no mortality. No pulmonary emboli seen.
Rogers (6)	1993	Class III	Retrospective review of 2525 high-risk trauma patients. Four high-risk groups that account for 92% of PE were identified. <ol style="list-style-type: none"> <li>1. Patients &gt; 55 years of age with isolated long bone fractures</li> <li>2. Patients with severe head injury and coma</li> <li>3. Patients with multiple long bone fractures and pelvic fracture</li> <li>4. Patients with spinal cord injury and paraplegia or quadriplegia</li> </ol> Overall incidence of PE was 1%.
Winchell (7)	1994	Class III	Retrospective review of 9721 trauma patients. 0.37% sustained a clinical or autopsy documented PE. Only 22% had a known DVT. 80% of patients with PE were receiving some form of prophylaxis (including 22% who were receiving both pneumatic compression stockings AND subcutaneous heparin). High-risk patient categories included <ol style="list-style-type: none"> <li>1. Head and spinal cord injury</li> <li>2. Head and long bone fracture</li> <li>3. Severe pelvis and long bone fracture</li> <li>4. Multiple long bone fractures</li> </ol>
Rosenthal (8)	1994	Class III	Retrospective case-control study of 151 trauma patients evaluating an aggressive approach to IVC filter placement in high-risk patients. From 1984-1988, 19 of 94 patients (20%) developed DVT despite prophylaxis (mechanical/ subcutaneous heparin). 8 patients developed PE (2 fatal). 15% of patients sustained PE without DVT (3 fatal). No patient sustained PE after filter placement. 23% of patients with ISS>16 developed PE.  From 1988-1992, 29 of 67 patients with ISS>16 had filters placed. 13% of all patients developed DVT. Only 1% of patients with ISS>16 developed PE with the more aggressive approach.
Wilson (9)	1994	Class III	Retrospective evaluation of PE in 2525 trauma patients. 6% of patients with traumatic spinal cord injury (SCI) developed PE. Following a more aggressive utilization of IVC filters, no PE has been noted in 15 patients with SCI over a 6-24 month follow-up period.
Khansarinia (10)	1995	Class II	Prospective case-control evaluation of prophylactic IVC filters in 224 patients. 0% incidence of PE in 108 patients with prophylactic IVC filter vs. 6% in 216 historically matched control patients (4% fatal) (p<0.009).
Rodriguez (11)	1996	Class III	Prospective case (40 patients) vs. injury-matched historical control (80 patients) study. PE decreased from 14% to 1% (p=0.02) with prophylactic IVC filter placement. 44% of PE's occurred in the first week.
Gosin (12)	1997	Class III	Prospective case (250 patients) vs. historical control (249 patients) study. Prophylactic IVC filter placement in high-risk trauma patients decreased the PE rate from 4.8% to 1.6% (p=0.045). No clinically evident complications of IVC filter placement were noted.
Rogers (13)	1997	Class II-III	Retrospective review of high-risk orthopedic trauma patients. High-risk injury patterns for PE included: <ol style="list-style-type: none"> <li>1) Lower extremity fractures (0.62%)</li> <li>2) Pelvic fractures (1.3%)</li> <li>3) Pelvic and LE fractures (2.5%)</li> <li>4) Non-orthopedic trauma patients (0.15%)</li> </ol> IVC filters were placed in 35 of 940 patients who met 2 or more of the following criteria: <ol style="list-style-type: none"> <li>1) Age &gt; 55 years</li> <li>2) ISS &gt; 16</li> <li>3) Complex pelvic fractures</li> <li>4) Long bone and pelvic fractures</li> <li>5) Lower extremity or pelvic fracture requiring prolonged bedrest</li> </ol>

			Incidence of PE decreased from historical rate of 1% to 0.2% in study population ( $p < 0.04$ ).
Rogers (14)	1998	Class II	Prospective evaluation of IVC filter placement in 792 trauma patients with 35 at high-risk and a contraindication to anticoagulation. No high-risk patient developed PE with a filter in place. 0.25% incidence of PE in trauma patients not deemed to be at high-risk.
Rajasekhar (28)	2011	Class III	Meta-analysis of 7 observational studies with total of 1,900 patients. Rate of PE was statistically lower in the IVC filter group compared to a matched control group without IVC filters. These results demonstrated decreased likelihood of PE among trauma patients who received filters. However, results not generalizable secondary to lack of contemporary use of pharmacologic prophylaxis across all studies.
Shariff (29)	2020	Class III	Meta-analysis of 10 studies, two of which were randomized controlled trials. Use of prophylactic IVC filters was associated with a reduced risk of symptomatic PE among patients with major trauma (ISS > 15), however the use of IVC filters was not associated with a decreased risk of fatal PE.
Kwok (30)	2019	Class I	Multicenter, randomized control trial. Enrolled 240 severely injured patients either to IVC filter group or no filter group. Found that early placement of a vena cava filter did not result in significantly lower incidence of symptomatic PE or death vs. no placement of a filter (13.9% IVC filter group vs. 14.4% control group). Also found an entrapped thrombus in in the filter of 6 patients. Concluded that early prophylactic placement of a vena cava filter did not result in lower incidence of symptomatic PE or death at 90 days vs. no filter.
Hemmila (31)	2015	Class III	Retrospective study comparing prophylactic IVC filter placement in 803 patients vs control set of 39,456 patients. Patients with IVC filter demonstrated no variation in mortality. Prophylactic IVC filter placement was associated with a statistically significant increase in incidence of DVT ( $p = 0.01$ ).

#### Bedside Insertion, Ultrasound Guidance and Temporary Filters

Author	Year	Evidence	Findings
Nunn (15)	1997	Class II	55 patients undergoing bedside IVC filter placement under ultrasound guidance. 89% had successful placement. Failures were mostly due to inability to visualize the right renal vein due to bowel gas. No procedure related mortality and no PE. Four complications (8.2%) included 1 tilted filter, 1 DVT at the needle puncture site, 1 IVC occlusion, and 1 minor filter migration. Estimated annual cost savings were significant (\$69,800-\$118,300). Various other reports confirm the safety, feasibility, and cost effectiveness of this approach (13-16).
Linsenmaier (16)	1998	Class II	Prospective evaluation of 50 temporary IVC filter (Gunther, Gunther Tulip, Antheor) placements. 100% placement success. All temporary filters were removed in 1-12 days (mean 7.3 days). On removal, 18% showed thrombi in the filter. No patients developed a PE with a filter in place. 2 filters migrated and 1 patient developed an IVC thrombosis. 2 filters required femoral venotomy for removal.
Offner (17)	2003	Class II	Prospective evaluation of 44 temporary IVC filter (Gunther Tulip) placements. 84% were in severely injured patients. Filters were in place an average of $14 \pm 1$ days (range 3-30 days). Three filters could not be retrieved, 2 because of significant clots below the filter and 1 because of abnormal angulation. No complications associated with insertion or retrieval.

### Timing of Prophylaxis

Author	Year	Evidence	Findings
Owings (18)	1997	Class III	Retrospective review of 63 trauma patients with PE. 25% of PE's occurred within the first 4 days of injury. 4 patients had their PE (1 fatal) the day following injury.
Carlin (19)	2002	Class III	Retrospective review of 22 trauma patients who developed PE prior to IVC filter placement. On average, PE was diagnosed $4 \pm 2$ days from admission and 36% occurred in the first 72 hours.

### Follow up & Complications

Author	Year	Evidence	Findings
Greenfield (20)	1995	Class III	20-year follow-up of long-term safety and efficacy of IVC filter placement. Data were available for 54% of placements. Mean follow-up was 56.5 months. 93% had a patent insertion site vein. 5% had significant tilting or migration. 2% had a fractured filter strut. No clinical sequelae were noted for tilt, migration or limb fracture. Caval patency was 96%.
Rogers (13)	1998	Class III	Retrospective review of prophylactic IVC filter placement in 132 trauma patients. 3% demonstrated insertion-related thrombosis and 2.3% PE. 36% had follow-up ultrasound examinations. Mean follow-up time 599 days (range 9-1946 days). One asymptomatic IVC thrombosis was detected. 5.5% demonstrated strut malpositioning with a higher incidence of PE in these patients (6.3% vs. 0%; $p=0.05$ ).
Langan (21)	1999	Class III	Retrospective review of 160 trauma patients with prophylactic IVC filters. 47% survey response rate and return for examination, duplex ultrasound, and fluoroscopy. Mean follow up was 19.4 months (range 3 to 57 months). The IVC was visualized in 93% and patency was 100% in these patients. Fluoroscopy failed to show any evidence of filter migration. One known clinical PE in 187 patients (0.5%) in whom a filter was inserted.
Sekharan (22)	2000	Class III	Retrospective review of 90 multi-system trauma patients receiving prophylactic IVC filters. 37% returned for evaluation. Mean follow-up was 68 months. 6% demonstrated DVT, 18% lower extremity edema, 0% PE, 0% migration / limb fracture. No IVC thrombosis.
Greenfield (23)	2000	Class III	Retrospective review of IVC filters in 385 trauma patients (249 prophylactic). Long-term outcome was available in 79%. Mean follow up 2.4 years. 2% had insertion site thrombosis and 15.6% DVT. Migration and tilt were rare and clinically and statistically insignificant. IVC patency was 96.5%. 3 PE's (1.5%).
Wojcik (24)	2000	Class III	Retrospective review of 178 trauma patients. 59% returned for follow-up. Mean follow-up 28.9 months. No clinically symptomatic pulmonary emboli. One IVC filter migration (0.95%). One IVC occlusion (0.95%). In the prophylactic group ( $n=64$ ), 28 (44%) developed a DVT. 11 patients (10.4%) had LE swelling.
Duperier (25)	2003	Class III	Retrospective review of 133 trauma patients receiving IVC filters. 77% had post-insertion duplex studies. 26% had de novo thrombi. No arteriovenous fistulae were noted. No patients developed clinical evidence of IVC occlusion. One patient had a fatal PE.
Prepic Study Group (26)	2005	Class I	Four hundred patients randomized to permanent IVC filter placement vs. no filter in addition to standard anticoagulation were reassessed 8 years post-study. Symptomatic PE occurred in 6.2% of the filter group and 15.1% in the no-filter group ( $p=0.008$ ). DVT occurred in 35.7% of the filter group and 27.5% of the no-filter group ( $p=0.042$ ). Post-thrombotic syndrome occurred equally between the groups. There was no difference in long-term mortality. The authors concluded that while IVC filters reduce the risk of PE, they increase the risk of DVT and do not alter mortality. The prophylactic insertion of such filters in the general population with DVT cannot be recommended.
Singh (27)	2008	Class III	Retrospective review of 558 patients receiving an IVC filter. 362 filters met currently accepted indications while 196 filters did not (i.e., did not

			have a contraindication to or had not failed anticoagulation). The within-guidelines group had a 1.4% post-filter PE incidence, a 13.6% IVC thrombosis rate, and 9.4% with DVT. The out of guidelines group had a 0.5% post-filter PE incidence, a 1% IVC thrombosis rate, and 3.6% with DVT. No patient without DVT at IVC filter insertion subsequently developed a PE. The authors concluded that IVC filter placement cannot be supported in patients without DVT who can be anticoagulated.
--	--	--	--

## REFERENCES

1. Rogers FB, Cipolle MD, Velmahos G, Rozycki G. Management of venous thromboembolism in trauma patients. [www.east.org](http://www.east.org).
2. Stevens SM, Woller SC, Baumann Kreuziger L, Vintch JRE, Wells PS, et al. Antithrombotic Therapy for VTE Disease: Second Update of the CHEST Guideline and Expert Panel Report. *Chest* 2021; 160:e545-3608.
3. *BEST PRACTICES GUIDELINES THE MANAGEMENT OF TRAUMATIC BRAIN INJURY*. (n.d.). American College of Surgery Trauma Programs. Retrieved October 30, 2024, from <https://www.facs.org/media/vgfgjpfk/best-practices-guidelines-traumatic-brain-injury.pdf>
4. Elkbuli A, Ehrhardt JD, Kinslow K, McKenney M. Prophylactic inferior vena cava filters: outcomes in severely injured trauma patients. *Am Surg* 2021; 87(2):300-308.
5. Leach TA, Pastena JA, Swan KG, Tikellis JI, Blackwood JM, Odom JW. Surgical prophylaxis for pulmonary embolism. *Am Surg* 1994; 60:292-295.
6. Rogers FB, Shackford SR, Wilson J, Ricci MA, Morris CS. Prophylactic vena cava filter insertion in severely injured trauma patients: indications and preliminary results. *J Trauma* 1993; 35:637-642.
7. Winchell RJ, Hoyt DB, Walsh JC, Simons RK, Eastman AB. Risk factors associated with pulmonary embolism despite routine prophylaxis: implications for improved protection. *J Trauma*, 1994;37:600-606.
8. Rosenthal D, McKinsey JF, Levy AM, Lamis PA, Clark MD. Use of the Greenfield filter in patients with major trauma. *Cardiovascular Surgery* 1994; 2:52-55.
9. Wilson JT, Rogers FB, Wald SL, Shackford SR, Ricci MA. Prophylactic vena cava filter insertion in patients with traumatic spinal cord injury: preliminary results. *Neurosurgery* 1994; 35 234-239.
10. Khansarinia S, Dennis J, Veldenz H, Butcher JL, Hartland L. Prophylactic Greenfield filter placement in selected high-risk trauma patients. *J Vasc Surg* 1995; 22:231-236.
11. Rodriguez JL, Lopez JM, Proctor MC, Conley JL, Gerndt SJ, et al. Early placement of prophylactic vena caval filters in injured patients at high risk for pulmonary embolism. *J Trauma* 1996; 40:797-804.
12. Gosin JS, Graham AM, Ciocca RG, Hammond JS. Efficacy of prophylactic vena cava filters in high-risk trauma patients. *Ann Vasc Surg* 1997; 11:100-105.
13. Rogers FB, Shackford SR, Ricci MA, Huber BM, Atkins T. Prophylactic vena cava filter insertion in selected high-risk orthopaedic trauma patients. *J Ortho Trauma* 1997; 11:267-72.
14. Rogers FB, Strindberg G, Shackford SR, Osler TM, Morris CS, et al. Five-year follow-up of prophylactic vena cava filters in high-risk trauma patients. *Arch Surg* 1998; 133:406-412.
15. Nunn CR, Neuzil D, Naslund T, Bass JG, Jenkins JM, et al. Cost effective method for bedside insertion of vena caval filters in trauma patients. *J Trauma* 1997; 43:752-58.
16. Linsenmaier U, Rieger J, Schenk F, Rock C, Mangel E, Pfeifer KJ. Indications, management, and complications of temporary inferior vena cava filters. *Cardiovascular and Interventional Radiology*, 1998; 21:464-469.
17. Offner PJ, Hawkes A, Madayag R, Seale F, Maines C. The role of temporary inferior vena cava filters in critically ill surgical patients. *Arch Surg*, 2003; 138:591-595.
18. Owings JT, Kraut E, Battistella F, Cornelius JT, O'Malley R. Timing of the occurrence of pulmonary embolism in trauma patients. *Arch Surg*, 1997; 132:862-867.
19. Carlin AM, Tyburski JG, Wilson RF, Steffes C. Prophylactic and therapeutic inferior vena cava filters to prevent pulmonary emboli in trauma patients. *Arch Surg*, 2002; 137:521-527.
20. Greenfield LJ, Proctor MC. Twenty-year experience with the Greenfield filter. *Cardiovascular Surgery*, 1995; 3:199-205.
21. Langan EM, Miller RS, Casey WJ, Carsten CG, Graham RM et al. Prophylactic inferior vena cava filters in trauma patients at high risk: Follow-up examination and risk/benefit assessment. *J Vasc Surg* 1999; 30:484-490.
22. Sekharan J, Dennis JW, Miranda EF, Hertz JA, Veldenz HC, et al. Long-term follow-up of prophylactic Greenfield filters in multisystem trauma patients. *J Trauma*, 2000; 51:1087-1091.
23. Greenfield LJ, Proctor MC, Michaels AJ, Taheri PF. Prophylactic vena caval filters in trauma: the rest of the story. *J Vasc Surg* 2000; 32:490-497.

24. Wojcik R, Cipolle MD, Fearen I, Jaffe J, Newcomb J et al. Long-term follow-up of trauma patients with a vena caval filter. *J Trauma*, 1999; 49:839-843.
25. Duperier T, Mosenthal A, Swan K, Kaul S. Acute complications associated with Greenfield filter insertion in high-risk trauma patients. *J Trauma* 2003; 54:545-549.
26. Prepic Study Group. Eight-year follow-up of patients with permanent vena cava filters in the prevention of pulmonary embolism: the PREPIC (Prevention du Risque d'Embolie Pulmonaire par Interruption Cave) randomized study. *Circulation*. 2005; 112(3):416-22.
27. Singh P, LAI HM, Lerner RG, Chugh T, Aronow WS. Guidelines and the use of inferior vena cava filters: a review of an institutional experience. *Journal of Thrombosis and Haemostasis* 2008; 7: 65–71.
28. Pulmonary embolism prophylaxis with inferior vena cava filters in trauma patients: a systematic review using the meta-analysis of observational studies in epidemiology (MOOSE) guidelines - Database of Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews - NCBI Bookshelf (nih.gov)
29. Shariff M, Kumar A, Adalja D, Doshi R. Inferior vena cava filters reduce symptomatic but not fatal pulmonary emboli after major trauma: a meta-analysis with trial sequential analysis. *Eur J Trauma Emerg Surg* 2021; 47(6):1805-1811.
30. Kwok MH, Rao S, Honeybul S, Zellweger R, Wibrow B, et al. A multicenter trial of vena cava filters in severely injured patients. *NEJM* 2019; 381:328-337.
31. Hemmila MR, Osborne NH, Henke PK, Kepros JP, Patel SG, et al. Prophylactic inferior vena cava filter placement does not result in a survival benefit for trauma patients. *Ann Surg* 2015; 262(4):577-585.